



Communicable Disease Prevention Daily Health Check Questionnaire

All visitors must review this important information

All individuals are required to perform a daily health check at home prior to attending /visiting this facility. The BC Centre for Disease Control has provided the following recommendations for daily health assessment.

Symptoms of illness and Screening Criteria

Are you experiencing any of following symptoms?

- | | |
|------------------------------------------------------------|-------------------------------------------------------|
| <input type="checkbox"/> Fever (>38.0° C) | <input type="checkbox"/> Diarrhea |
| <input type="checkbox"/> Chills | <input type="checkbox"/> Headache |
| <input type="checkbox"/> New or worsening of chronic cough | <input type="checkbox"/> Sore throat |
| <input type="checkbox"/> Loss of sense of smell or taste | <input type="checkbox"/> Loss of appetite |
| <input type="checkbox"/> Difficulty breathing | <input type="checkbox"/> Extreme fatigue or tiredness |
| <input type="checkbox"/> Nausea or vomiting | <input type="checkbox"/> Body ache |

- **Tested positive for COVID-19** (< 14 days) or currently being investigated for COVID-19?
- **Self-Isolation:** You or anyone in your household been told to **be in self-isolation now?**
- **Are you a confirmed close contact** of a person confirmed to have COVID-19?

Based on the information above, if you answered “yes” to any of the above, do not enter this site, leave the ground immediately and contact 811 for guidance.

Also, **DO NOT ENTER** if you have travelled internationally in the last 14 days and are required to be in isolation.

DISTRICT PROTOCOLS



Please follow all safety protocols: Wash your hands, respect personal space and wear a mask.

If you develop COVID-19 like symptoms while on the premises, follow the following steps: during your workday follow the following steps:

1. Alert the site Administrator
2. Go home and self-isolate
3. Monitor your symptoms
4. Get assessed by healthcare provider. Testing for COVID-19 is recommended, even if mild COVID-19 symptoms.