## **BC ROUTINE IMMUNIZATION SCHEDULE**

## **INFANTS & CHILDREN**

|  | CHILD'S AGE  |             |                    |              |              |   |
|--|--------------|-------------|--------------------|--------------|--------------|---|
| VACCINE  | 2<br>months  | 4<br>months | 6<br>months        | 12<br>months | 18<br>months | Starting at 4<br>years<br>(kindergarten<br>entry) |
| DTaP-HB-IPV-Hib<br>(diphtheria, tetanus, pertussis, hepatitis B, polio,<br><i>Haemophilus influenzae</i> type b)<br>HealthLinkBC File # <u>105</u> | ~            | 1           | <b>√</b>           |              |              |   |
| Pneumococcal conjugate <sup>‡</sup><br>HealthLinkBC File # <u>62a</u>  | 1            | <b>√</b>    |                    | 1            |              |   |
| Rotavirus<br>HealthLinkBC File # <u>104</u>  | $\checkmark$ | 1           |                    |              |              |   |
| Meningococcal conjugate C<br>HealthLinkBC File # 23a   | 1            |             |                    | <b>√</b>     |              |   |
| MMR<br>(measles, mumps, rubella)<br>HealthLinkBC File # <u>14a</u>   |              |             |                    | <b>√</b>     |              |   |
| Varicella <sup>1</sup><br>(chickenpox)<br>HealthLinkBC File # <u>44b</u>   |              |             |                    | 1            |              |   |
| DTaP-IPV-Hib<br>(diphtheria, tetanus, pertussis, polio,<br><i>Haemophilus influenza</i> e type b)<br>HealthLinkBC File # <u>15b</u>                |              |             |                    |              | $\checkmark$ |   |
| DTaP-IPV<br>(diphtheria, tetanus, pertussis, polio)<br>HealthLinkBC File # <u>15d</u>  |              |             |                    |              |              | 1   |
| MMRV <sup>1</sup><br>(measles, mumps, rubella, varicella)<br>HealthLinBC File # <u>14e</u>   |              |             |                    |              |              | <b>√</b>  |
| Influenza<br>HealthLinkBC File # <u>12d</u> and <u>12e</u>   |              |             | (6 months-4 years) |              |              |   |
| Hepatitis A <sup>+</sup><br>HealthLinkBC File # <u>33</u>  |              |             | $\checkmark$       |              |              | <b>v</b> .  |

Children with specific medical conditions that place them at high risk of disease should receive an additional dose at 6 months of age. See HealthLinkBC File # 62a for more information or speak to your health care provider.

- Children who had chickenpox or shingles disease at 1 year of age or older do not need the chickenpox vaccine. Children entering school who need both a 2<sup>nd</sup> dose of MMR and varicella vaccines may be immunized with MMRV vaccine.
- \* A second dose is needed 4 weeks after the first dose if receiving vaccine for the first time.
- Hepatitis A vaccine is offered to Aboriginal children living both on-reserve and off-reserve.
- Hepatitis A vaccine is offered to Aboriginal children starting at 6 months of age. Alternatively, 1 or 2 doses will be offered to kindergarten-age children starting at 4 years of age depending on previous immunization history.